

RESCUE®

READY TO USE

ONE STEP DISINFECTANT CLEANER & DEODORIZER

For Veterinary Use

Reference Sheet

**KEEP OUT OF REACH
OF CHILDREN**

Manufactured by:

Virox Technologies Inc.

2770 Coventry Road
Oakville, ON Canada L6H 6R1
viroxanimalhealth.com



EPA Est. No. 74559-CAN-1
EPA Reg. No. 74559-9

This product is a one-step germicidal disinfectant cleaner and deodorant designed for general cleaning and disinfecting of hard, non-porous environmental surfaces.

It cleans quickly by removing dirt, grime, blood, urine, mold stains, laboratory stains, and other common soils found in animal housing facilities, grooming facilities, kennels, pet stores, veterinary clinics, or other small animal facilities.

This product is designed for use on hard, non-porous environmental surfaces: chrome, glazed ceramic, glazed porcelain, glazed tile, stainless steel, associated with high touch environmental surfaces, animal equipment, animal crates, cages, carts, examination tables, grooming equipment, reception areas, veterinary care surfaces or any hard non-porous washable (non-food contact) surfaces where disinfection is required. It also reduces odor-causing bacteria leaving surfaces smelling clean and fresh.

BACTERICIDAL IN 1 MINUTE

Staphylococcus aureus
Staphylococcus aureus (Methicillin Resistant (MRSA))
Salmonella enterica (formerly known as *Salmonella choleraesuis*)
Klebsiella pneumoniae
Klebsiella pneumoniae (Extended Spectrum Beta-lactamase (ESBL))
Klebsiella pneumoniae (New Delhi metallo-β-lactamase 1 (NDM-1))
Klebsiella pneumoniae (Carbapenem Resistant (CRE))
Staphylococcus aureus (Vancomycin Intermediate Resistant (VISA))
Staphylococcus aureus (Vancomycin Resistant (VRSA))
Staphylococcus pseudintermedius
Staphylococcus epidermidis (Methicillin Resistant (MRSE))
Enterococcus faecalis (Vancomycin Resistant (VRE))
Enterococcus faecium
Bordetella bronchiseptica
Streptococcus pneumoniae (Penicillin Resistant (PRSP))
Pasteurella multocida
Serratia marcescens
Shigella dysenteriae

*VIRUCIDAL IN 30 SECONDS

Adenovirus Type 8
Avian Influenza A Virus (H7N9)
Avian Influenza A Virus (H3N2)
Cytomegalovirus
Hepatitis B Virus (HBV)
Hepatitis C Virus (HCV)
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2
HIV Type 1 Strain HTLV III_B
Human Corona Virus
Influenza A Virus (Hong Kong Strain)
Influenza B Virus
Norovirus (Feline Calicivirus as the surrogate)
Pandemic Influenza A Virus H1N1
Parainfluenza Type 3
Polio Type 1 (Strain Brunhilde)
Respiratory Syncytial Virus (RSV)

Rhinovirus R37
Rotavirus (WA strain)
Vaccinia Virus

ANIMAL PREMISE *VIRUCIDAL ACTIVITY IN 1 MINUTE, UNLESS OTHERWISE NOTED

Canine Parvovirus
Bovine Viral Diarrhea Virus – 30 seconds
Duck Hepatitis B virus – 30 seconds
Equine Influenza A Virus (H3N8) (EIV) – 30 seconds
Feline Calicivirus- 30 seconds
Feline Infectious Peritonitis Virus (FIPV)
Feline Herpes Virus (Feline Rhinotracheitis)
Murine Hepatitis Virus
Minute Virus of Mouse (Murine Parvovirus)
Murine Norovirus – 30 seconds
Feline Panleukopenia Virus
Canine Distemper Virus

This product has demonstrated effectiveness against Influenza Type A₂ (Hong Kong), and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

FUNGICIDAL

Trichophyton mentagrophytes
Candida albicans – 1 minute
Microsporum canis – 1 minute

MALODORS

Malodors Activity – eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

MOLD/MILDEW

Mold/Mildew (Mildewcidal Activity) – kills the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew (Mildewstatic Activity) – controls and prevents the growth of mold and mildew at a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

TUBERCULOCIDAL IN 1 MINUTE

Mycobacterium bovis (BCG)

SOFT SURFACE SANITIZATION AGAINST THE FOLLOWING ORGANISMS IN 30 SECONDS:

Staphylococcus aureus

Enterobacter aerogenes

Methicillin Resistant *Staphylococcus aureus* (MRSA)

Vancomycin Resistant *Enterococcus faecalis* (VRE)

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER IN 10 SECONDS AGAINST:

Staphylococcus aureus

Enterobacter aerogenes

Acinetobacter baumannii

Listeria monocytogenes

Campylobacter jejuni

Salmonella enterica (formerly known as *Salmonella choleraesuis*)

Pseudomonas aeruginosa

Escherichia coli O157:H7

Vancomycin Resistant *Enterococcus faecalis* (VRE)

Klebsiella pneumoniae

Methicillin Resistant *Staphylococcus aureus* (MRSA)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6-8 inches from surface.
3. Allow surface to remain wet for 1 minute.
4. Wipe surfaces dry or allow to air dry.

To Disinfect Non-Critical[¥], Pre-Cleaned Instruments:

1. Instruments must be thoroughly pre-cleaned to remove excess organic debris, rinsed, and rough dried.
2. Clean and rinse lumens of hollow instruments before spraying with this product.
3. Spray all surfaces of instruments with this product until thoroughly wet.
4. Let stand for 1 minute.
5. Wipe with a clean, damp cloth or paper towel and allow to air dry.

¥ Non-Critical medical devices are items that come in contact only with intact skin.

For use as sanitizer on Soft Surfaces or Fabrics[‡]:

1. Hold bottle upright 6"-8" from surfaces.
2. Spray until fabric is wet. Do not saturate.
3. Let stand for 30 seconds. Allow to air dry.
4. For difficult odors or heavy fabrics, repeat application. Reapply as necessary.

‡ composed of cotton or polyester

For Use as a Cleaner and/or Deodorizer:

Spray 6-8 inches from non-porous environmental surfaces. Allow to air dry.

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood / body fluids, and in which the surfaces / objects likely to be soiled with blood / body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS Virus), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood / body fluids must be thoroughly cleaned from surfaces / objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV.

Disposal of Infectious Material: Blood / body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Kill Fungi: Pre-clean heavily soiled areas. Spray product directly onto non-porous surfaces until completely wet. Allow surface to remain wet for 30 seconds. Wipe surfaces and let air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Spray product directly onto non-porous surfaces until completely wet. Allow the surface to remain visibly wet for 4 minutes. Allow to air dry or wipe surface dry.

To Control Mold and Mildew: Spray product directly onto non-porous surfaces until completely wet. Allow to air dry or wipe surface dry.

Animal Housing Facilities in Veterinary Clinics Hospitals, Farm Premise Including, Equine production Farm, Bovine production Farm:

For cleaning and disinfecting the following hard non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities including barns, pens, stalls, chutes, and barns occupied or traversed by animals.
3. Empty (or cover) all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all surfaces floors, walls, cages and other washable hard, non-porous environmental surfaces by spraying with solution until (thoroughly) wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. For disinfection of feed racks, troughs, automatic feeders, fountains and water appliances waterers spray with solution until completely wet and let stand for 1 minute. Then thoroughly scrub all treated surfaces with soap or detergent, and rinse with potable water before reuse.

Allergen Destruction

To destroy specified allergens: Spray, wait 1 minute, and wipe excess. Allow to air dry.

Allergens:

Cockroach matter -or- particles

Dust mite matter -or- particles

Pet dander -or- dog and cat dander